

# Sequential Convex Approximations to Joint Chance Constrained Programs: A Monte Carlo Approach

L. Jeff Hong

Department of Industrial Engineering and Logistics Management, The Hong Kong University of Science and Technology,  
Clear Water Bay, Hong Kong, China, hongl@ust.hk

Yi Yang

Department of Computer Science, University of California, Irvine, California 92617, yangyi02@gmail.com

Liwei Zhang

School of Mathematical Science, Dalian University of Technology, Dalian 116024, China, lwzhang@dlut.edu.cn

When there is parameter uncertainty in the constraints of a convex optimization problem, it is natural to formulate the problem as a joint chance constrained program (JCCP), which requires that all constraints be satisfied simultaneously with a given large probability. In this paper, we propose to solve the JCCP by a sequence of convex approximations. We show that the solutions of the sequence of approximations converge to a Karush-Kuhn-Tucker (KKT) point of the JCCP under a certain asymptotic regime. Furthermore, we propose to use a gradient-based Monte Carlo method to solve the sequence of convex approximations.

*Subject classifications:* programming, stochastic: chance constrained program.

*Area of review:* Simulation.

*History:* Received March 2009; revisions received June 2009, December 2009, January 2010; accepted February 2010.

## 1. Introduction

Consider the following optimization problem:

$$\begin{aligned} & \text{minimize} && h(x), \\ & \text{subject to} && c_1(x, \xi) \leq 0, \dots, c_m(x, \xi) \leq 0, \\ & && x \in X, \end{aligned} \quad (1)$$

where  $\xi$  is a  $k$ -dimensional parameter vector,  $X$  is a subset of  $\mathfrak{R}^d$ ,  $h: \mathfrak{R}^d \rightarrow \mathfrak{R}$  and  $c_i: \mathfrak{R}^{d+k} \rightarrow \mathfrak{R}$ ,  $i = 1, \dots, m$ , are real-valued functions. Furthermore, we assume that  $h(x)$  and  $c_i(x, \xi)$ ,  $i = 1, \dots, m$  are convex in  $x$  and  $X$  is a compact convex set. Then, Problem (1) is a standard constrained convex optimization problem. It has broad applications in communications and networks, product design, system control, statistics, and finance, and it can be solved efficiently (e.g., see Boyd and Vandenberghe 2004 for a comprehensive introduction to convex optimization).

In many practical problems, however, the parameter vector  $\xi$  of Problem (1) might be uncertain. If this uncertainty is ignored (e.g., by using the expected values of  $\xi$  in the optimization), the optimal solution obtained by solving Problem (1) might actually be infeasible with a very high probability. To illustrate this, we consider a very simple example. We let  $\xi = (\xi_1, \dots, \xi_m)$  where  $\xi_1, \dots, \xi_m$  are  $m$  independent observations of a standard normal distribution, let  $X = \mathfrak{R}$ ,  $h(x) = x$ , and  $c_i(x, \xi) = \xi_i - x$  for all  $i = 1, \dots, m$ . If we ignore the parameter uncertainty by using  $E(\xi)$  to substitute  $\xi$  in Problem (1), the optimal solution is

$x^* = 0$ . However, the probability of  $x^* = 0$  being a feasible solution equals

$$\begin{aligned} & \Pr\{c_1(x^*, \xi) \leq 0, \dots, c_m(x^*, \xi) \leq 0\} \\ & = \Pr\{x^* \geq \xi_1\} \dots \Pr\{x^* \geq \xi_m\} = 0.5^m, \end{aligned}$$

which is very small when  $m$  is large (for instance, it is already less than 0.001 when  $m = 10$ ).

To consider this parameter uncertainty, we may formulate the problem as

$$\begin{aligned} \text{(P)} \quad & \text{minimize} && h(x), \\ & \text{subject to} && \Pr\{c_1(x, \xi) \leq 0, \dots, c_m(x, \xi) \leq 0\} \geq 1 - \alpha, \\ & && x \in X. \end{aligned}$$

In Problem (P), we require all  $m$  uncertain constraints be satisfied simultaneously with a probability at least  $1 - \alpha$ , where  $0 < \alpha < 1$  is often set as 0.01, 0.05, or 0.1. Therefore, the solution to Problem (P) is guaranteed to be a feasible solution to the original Problem (1) with a probability at least  $1 - \alpha$ . Problem (P) is called a joint chance constrained program (JCCP), and the probabilistic constraint is called a joint chance constraint. When  $m = 1$ , the constraint is called a single chance constraint because it requires only a single constraint to be satisfied with probability  $1 - \alpha$ . For simplicity of the notation, we let

$$p(x) = 1 - \Pr\{c_1(x, \xi) \leq 0, \dots, c_m(x, \xi) \leq 0\},$$

and  $p(x)$  is the probability that at least a constraint is violated. Then, the joint chance constraint of Problem (P) becomes  $p(x) \leq \alpha$ .

Many stochastic optimization problems can be formulated as a JCCP. For instance, the reservoir system design problem of Prékopa et al. (1978) minimizes the total building and penalty costs while satisfying demands for all sites and all periods with a probability at least 80%, and the cash matching problem of Dentcheva et al. (2004) maximizes the value of the portfolio at the end of the planning horizon while covering all scheduled payments with a probability at least 95%. JCCPs were first introduced and studied by Charnes et al. (1958), Miller and Wagner (1965), and Prékopa (1970). Since then, JCCPs have been studied extensively in the stochastic optimization literature. For a recent review of the topic, readers are referred to Prékopa (2003).

There are generally two major difficulties in solving a JCCP. First,  $p(x)$  might not be a convex (or quasiconvex) function even though  $c_1(x, \xi), \dots, c_m(x, \xi)$  are all convex in  $x$ . Therefore, Problem (P) might not be a convex optimization problem even though Problem (1) is. Then, it is difficult to find a global optimal solution. Second,  $p(x)$  generally has no closed form and is typically difficult to evaluate.

Different approaches have been proposed in the stochastic optimization literature to address these difficulties. For the convexity of Problem (P), Prékopa (2003) showed that  $p(x)$  is quasiconvex (which implies that Problem (P) is convex) if  $c_1(x, \xi), \dots, c_m(x, \xi)$  are quasiconvex functions of  $(x, \xi)$ , and if  $\xi$  has a logconcave probability distribution, which includes uniform distribution, multivariate normal distribution, and many others. Lagoa et al. (2005) showed that an individual chance constraint in the form of  $\Pr\{a^T x \leq b\} \geq 1 - \alpha$  defines a convex set provided that the vector  $(a^T, b)^T$  has a symmetric logconcave density with  $\alpha < 1/2$ . Henrion (2007) showed that an individual chance constraint in the form of  $\Pr\{\xi^T q(x) \leq b\} \geq 1 - \alpha$  defines a convex set provided that all components of  $q(x)$  are nonnegative and convex,  $\alpha < 1/2$ , and the vector  $\xi$  has an elliptically symmetric distribution whose parameters satisfy certain requirements. Henrion and Strugarek (2008) showed that a joint chance constraint in the form of  $\Pr\{g_i(x) \geq \xi_i, i = 1, \dots, m\} \geq 1 - \alpha$  defines a convex set if  $g_i(x)$  is  $(-r_i)$ -concave and  $\xi_i, i = 1, \dots, m$  are independent random variables with  $(r_i + 1)$ -decreasing densities for some  $r_i > 0$  for sufficiently small  $\alpha$  values.

When  $p(x)$  is not quasiconvex (or at least not verifiable), many convex conservative approximations of  $p(x)$  have been proposed, e.g., the quadratic approximation of Ben-Tal and Nemirovski (2000), the conditional value-at-risk (CVaR) approximation of Rockafellar and Uryasev (2000), and the Bernstein approximation of Nemirovski and Shapiro (2006). These approximations typically find feasible but suboptimal solutions to Problem (P). Furthermore, most of these approximations work only on single chance

constraints instead of a joint chance constraint. Therefore, one has to approximate the joint chance constraint by a set of individual chance constraints. A popular choice is to use Boole's inequality, which guarantees the satisfaction of the joint chance constraint if  $\Pr\{c_i(x, \xi) \geq 0\} \geq 1 - \alpha_i, i = 1, \dots, m$ , and  $\alpha_1 + \dots + \alpha_m = \alpha$  (e.g., Nemirovski and Shapiro 2006). However, it makes the solution even more conservative.

To evaluate  $p(x)$ , Monte Carlo simulations are often used when the closed form of  $p(x)$  is not available. When the chance constraint is approximated by functions that are analytically tractable, e.g., the quadratic approximation or the Bernstein approximation, evaluations of these functions are easy. The resulting problems can be solved directly using standard nonlinear optimization algorithms. When the chance constraint is approximated by functions that are not analytically tractable, e.g., the CVaR approximation, Monte Carlo simulations are also used to evaluate these functions (Rockafellar and Uryasev 2000).

Luedtke and Ahmed (2008) studied the sample-average approximation of the JCCPs. Their goal is to determine the sample size and appropriate probability requirement such that one can find a feasible solution of the original JCCP and also bound the optimality gap. Luedtke et al. (2007) considered linear programs with joint chance constraints and showed that the problems can be reformulated into mixed integer programs. They further demonstrated that the mixed integer programs can be solved efficiently when only the right-hand-side vector is random.

Another approach to solving JCCPs is to use the scenario approach, which solves the following problem:

$$\begin{aligned} & \text{minimize} && h(x), \\ & \text{subject to} && c_l(x, \xi_l) \leq 0, \quad i = 1, \dots, m, l = 1, \dots, n, \\ & && x \in X, \end{aligned} \quad (2)$$

where  $\xi_1, \xi_2, \dots, \xi_n$  are independent observations of  $\xi$  that are often generated from a Monte Carlo simulation. Problem (2) is a convex optimization problem and analytically tractable. The critical issue is how to determine the sample size  $n$  to ensure that the joint chance constraint is satisfied with a high probability. Calafiore and Campi (2005, 2006) and de Farias and Van Roy (2004) studied this issue independently, and Erdgoğan and Iyengar (2006) further extended the earlier results to situations where the distribution of  $\xi$  is ambiguous. The scenario approach is simple to understand and easy to implement. However, it also has several drawbacks. First, Problem (2) is also a conservative approximation to the original JCCP; it finds feasible but suboptimal solutions. Second, the solutions found by the scenario approach are not stable. They can be drastically different when different sets of samples are used. Third, the performance of the approach cannot be improved by acquiring more samples of  $\xi$ , which is in contrast to many other Monte Carlo algorithms. Indeed, increasing sample

size will make Problem (2) more conservative and might lead to worse solutions. Therefore, the performance of the approach cannot be improved when an ample amount of computational time is available.

In this paper, we propose a new approach to solving Problem (P). We first show that the function  $p(x)$  can be represented as a limit of a DC function (i.e., difference of two convex functions). Then we use an  $\varepsilon$ -approximation to approximate Problem (P). We show that, as  $\varepsilon$  goes down to zero, the optimal solutions (either global optimal or KKT points) of the approximation converge to the optimal solutions of Problem (P), respectively. To solve the  $\varepsilon$ -approximation problem, we propose to solve a sequence of convex optimization problems. We show that the sequence of solutions converge to a KKT point of the  $\varepsilon$ -approximation problem under moderate conditions. We also propose to use a Monte Carlo method to solve the sequence of convex optimization problems. We show that the solution of the Monte Carlo method converges with probability 1 as the sample size goes to infinity, and the sample problem can be solved efficiently using a gradient approach.

Compared to other approaches in the literature, our approach has several advantages. First, it converges to a KKT point of Problem (P), while nearly all other methods are conservative approximations whose solutions do not satisfy any optimality conditions of Problem (P). When the JCCP is convex (even though it might not be verifiable), our approach converges to the global optimal solution while other methods cannot. Second, our approach directly handles the joint chance constraint without breaking it into multiple single chance constraints. Therefore, it avoids the extra conservativeness introduced by using Boole's inequality.

Our approach also has several drawbacks compared to other approaches in the literature. First, it is computationally slow. Generally, Monte Carlo methods are slower than deterministic approximations because the sample problem is either of a large size or the functions need to estimate repeatedly. Our approach requires to solve a sequence of sample problems. Therefore, it can solve only problems with a small or moderate size, e.g., problems with less than 100 dimensions. Second, our approach requires the full joint distribution of the uncertain parameters in order to generate Monte Carlo samples. However, specifying a full joint distribution for a large number of parameters is often difficult in practice. When the distribution of  $\xi$  is ambiguous, our approach cannot be applied. To solve JCCPs with ambiguous distributions, many robust optimization approaches have been proposed in the literature, e.g., Ben-Tal and Nemirovski (2000), Bertsimas and Sim (2004), and Chen et al. (2010).

The rest of the paper is organized as follows. We provide a new formulation of the JCCP in §2, and we show how the formulation can be solved by sequential convex approximations in §3. In §4, we propose an efficient Monte

Carlo algorithm to solve the sequence of convex approximations. The numerical results are reported in §5, followed by the conclusions and future research in §6. An electronic companion to this paper is available as part of the online version that can be found at <http://or.journal.informs.org>. Some lengthy proofs are included in the electronic companion to this paper.

## 2. A DC Formulation

Let  $c(x, \xi) = \max\{c_1(x, \xi), \dots, c_m(x, \xi)\}$ . Note that  $c(x, \xi)$  is a convex function of  $x$  because  $c_i(x, \xi)$ ,  $i = 1, \dots, m$ , are all convex in  $x$ . Then,

$$\begin{aligned} p(x) &= 1 - \Pr\{c_1(x, \xi) \leq 0, \dots, c_m(x, \xi) \leq 0\} \\ &= 1 - \Pr\{c(x, \xi) \leq 0\} = \Pr\{c(x, \xi) > 0\}. \end{aligned}$$

By doing so, we convert a joint chance constraint to a single chance constraint. We can do this in this paper because we do not need to exploit certain special structures of  $c_i(x, \xi)$ . For many papers that require  $c_i(x, \xi)$  be a linear function of  $x$  or  $\xi$  and exploit this linearity, e.g., Ben-Tal and Nemirovski (2000) and Nemirovski and Shapiro (2006),  $c(x, \xi)$  is no longer linear when  $m > 1$ , and thus handling a joint chance constraint is significantly more difficult than handling a single chance constraint. For us, however, handling a joint chance constraint is as difficult as handling a single chance constraint.

A major difficulty of solving Problem (P) is that  $p(x)$  is generally not a convex function of  $x$  even though  $c(x, \xi)$  is convex in  $x$ . In the literature, many algorithms have been proposed to approximate  $p(x)$  by a conservative function  $\tilde{p}(x)$ , i.e.,  $\tilde{p}(x) \geq p(x)$  for all  $x \in X$ . Then, the solution to minimize  $h(x)$ ,

$$\text{subject to } \tilde{p}(x) \leq \alpha, \quad x \in X$$


is a feasible solution of Problem (P). If  $\tilde{p}(x)$  is close to  $p(x)$ , then the solution is a good approximation to the optimal solution of Problem (P). If  $\tilde{p}(x)$  is a convex function, then the approximated problem is a convex program that might be easier to solve.

In this section, we first introduce the CVaR approximation of Rockafellar and Uryasev (2000), which is the “best” convex conservative approximation (Nemirovski and Shapiro 2006). Based on the CVaR approximation, we propose another conservative approximation to  $p(x)$ , called the DC approximation. We then study the properties of the DC approximation.

### 2.1. CVaR Approximation

Note that  $p(x) = \Pr\{c(x, \xi) > 0\} = E[1_{(0, +\infty)}(c(x, \xi))]$ , where  $1_A(z)$  denotes the indicator function of set  $A$  that equals to 1 if  $z \in A$  and 0 if  $z \notin A$ . Because the indicator function  $1_{(0, +\infty)}(z)$  is nonconvex (see the left panel of Figure 1), one way to approximate  $p(x)$  is to find a convex approximation  $\psi(z)$  of  $1_{(0, +\infty)}(z)$  such that  $\psi(z) \geq 1_{(0, +\infty)}(z)$  for any  $z \in \mathfrak{R}$ . Then,  $\tilde{p}(x) = E[\psi(c(x, \xi))]$  is a

**Figure 1.** The CVaR approximation to the indicator function  $1_{(0,+\infty)}(z)$ .



convex conservative approximation of  $p(x)$ . For instance, both the CVaR approximation of Rockafellar and Uryasev (2000) and the Bernstein approximation of Nemirovski and Shapiro (2006) use this approach.

Among all convex conservative approximations of this kind, the CVaR approximation is known to be the “best” (Nemirovski and Shapiro 2006). It uses

$$\psi(z, t) = \frac{1}{t}[t+z]^+$$

to approximate  $1_{(0,+\infty)}(z)$ , where  $t > 0$  and  $a^+ = \max\{a, 0\}$  (see the right-hand panel of Figure 1), and let


$$p'(x) = \inf_{t>0} E[\psi(c(x, \xi), t)] = \inf_{t>0} \frac{1}{t} E[[t+c(x, \xi)]^+].$$

It can be shown that the new constraint  $p'(x) \leq \alpha$  is equivalent to  $\text{CVaR}_{1-\alpha}(c(x, \xi)) \leq 0$  (Nemirovski and Shapiro 2006), where

$$\text{CVaR}_{1-\alpha}(z) = \inf_{\tau \in \mathfrak{R}} \left\{ \tau + \frac{1}{\alpha} E[[z-\tau]^+] \right\}.$$

This is why this approximation is called the CVaR approximation. The CVaR approximation problem can be solved using a Monte Carlo method. Rockafellar and Uryasev (2000) provided an approach that solves the problem with a single chance constraint, and Nemirovski and Shapiro (2006) showed that the Boole’s inequality can be used to extend it to JCCPs. Hong and Liu (2009) provided a gradient-based Monte Carlo algorithm that directly solves the CVaR approximations of JCCPs.

**Figure 2.** The DC approximation to the indicator function  $1_{(0,+\infty)}(z)$ .



**2.2. DC Approximation**

Although the CVaR approximation may be the “best” convex conservative approximation of  $p(x)$ , it is clear that  $\psi(z, t)$  is not a good approximation to the indicator function  $1_{(0,+\infty)}(z)$  from Figure 1. The difference between the two functions grows unboundedly as  $z \rightarrow +\infty$ .

To find a better approximation to  $p(x)$ , we first define

$$\phi(z, t) = \frac{1}{t}[z]^+$$

for any  $t > 0$ . Note that  $\phi(z, t)$  can be obtained by shifting  $\psi(z, t)$  to the right side by a distance of  $t$  (see the left panel of Figure 2). Then,  $\pi(z, t) = \psi(z, t) - \phi(z, t)$  is a better approximation of  $1_{(0,+\infty)}(z)$  than  $\psi(z, t)$  is (see the right panel of Figure 2). Because both  $\psi(z, t)$  and  $\phi(z, t)$  are convex functions of  $z$ ,  $\pi(z, t)$  is a DC function of  $z$ . Furthermore, because  $\pi(z, t) \geq 1_{(0,+\infty)}(z)$  for all  $z \in \mathfrak{R}$  when  $t > 0$ ,  $\pi(z, t)$  is also a conservative approximation of  $1_{(0,+\infty)}(z)$  when  $t > 0$ .

Let  $g_1(x, t) = E[t+c(x, \xi)]^+$  and  $g_2(x) = g_1(x, 0)$ . Note that both functions are convex in  $x$ . Let

$$\tilde{p}(x, t) = E[\pi(c(x, \xi), t)] = \frac{1}{t}[g_1(x, t) - g_2(x)]. \tag{3}$$

Then,  $\tilde{p}(x, t)$  is a conservative DC approximation of  $p(x)$  for any  $t > 0$ . Let

$$\tilde{p}(x) = \inf_{t>0} \tilde{p}(x, t). \tag{4}$$

Then,  $\tilde{p}(x)$  is the best conservative approximation among all  $\tilde{p}(x, t)$  when  $t > 0$ . In this paper, we suggest to solve

$$(DC) \quad \begin{aligned} & \text{minimize} && h(x), \\ & \text{subject to} && \tilde{p}(x) \leq \alpha, \quad x \in X. \end{aligned}$$

We call Problem (DC) as the DC approximation of Problem (P).

### 2.3. Equivalence of the JCCP and DC Approximation

In this subsection, we prove that Problem (DC), which is a conservative approximation of Problem (P), is indeed equivalent to Problem (P). We make the following assumptions.

**ASSUMPTION 1.** *The set  $X$  is a compact and convex subset of  $\mathfrak{R}^d$ , and the support of  $\xi$ , denoted as  $\Xi$ , is a closed subset of  $\mathfrak{R}^k$ . For any  $\xi \in \Xi$ ,  $h(x)$ , and  $c_i(x, \xi)$ ,  $i = 1, \dots, m$ , are continuously differentiable and convex in  $x$  for any  $x \in \mathcal{O}$  where  $\mathcal{O}$  is a bounded open set, such that  $X \subset \mathcal{O}$ .*

Assumption 1 is used to clearly define Problem (P).

**ASSUMPTION 2.** *There exists a random variable  $K_i$  with  $E(K_i) < +\infty$ , such that*

$$|c_i(x_1, \xi) - c_i(x_2, \xi)| \leq K_i \|x_1 - x_2\|$$

for any  $x_1, x_2 \in \mathcal{O}$  and any  $i = 1, 2, \dots, m$ .

Let  $K = \sum_{i=1}^m K_i$ . Assumption 2 implies that  $|c(x_1, \xi) - c(x_2, \xi)| \leq K \|x_1 - x_2\|$  and  $E(K) < +\infty$ . The Lipschitz continuity of  $c(x, \xi)$  is critical in the analysis of differentiability of  $E[c(x, \xi)]$ . It is a common assumption used to handle the differentiability of an expectation (e.g., Broadie and Glasserman 1996, Hong 2009, and Hong and Liu 2010).

**ASSUMPTION 3.** *For any  $x \in \mathcal{O}$ ,  $c(x, \xi)$  is differentiable with respect to  $x$  w.p.1.*

To verify Assumption 3, we consider only the situation where  $\xi$  is a continuous random vector, because the following Assumption 4 is typically violated if  $\xi$  is discrete. By Assumption 1,  $c_i(x, \xi)$  is continuously differentiable for all  $i = 1, \dots, m$ . Therefore, if  $\Pr\{c_i(x, \xi) = c_j(x, \xi)\} = 0$  for any  $x \in \mathcal{O}$  and any  $i, j = 1, \dots, m$  with  $i \neq j$ ,  $c(x, \xi)$  is differentiable with respect to  $x$  w.p.1. In the electronic companion to this paper, we discuss how to verify and satisfy Assumption 3 if  $\Pr\{c_i(x, \xi) = c_j(x, \xi)\} \neq 0$ .

Let  $F(t, x) = \Pr\{c(x, \xi) \leq t\}$  be the cumulative distribution function of  $c(x, \xi)$ . We make the following assumption on the continuity of  $F(t, x)$ .

**ASSUMPTION 4.** *There exists a certain  $\delta > 0$  such that  $F(t, x)$  is continuously differentiable when  $(t, x) \in (-\delta, +\delta) \times \mathcal{O}$ .*

Because  $p(x) = 1 - F(0, x)$ , Assumption 4 implies that  $p(x)$  is continuously differentiable. Furthermore, note that  $\partial_t F(t, x)$  is the density function of  $c(x, \xi)$ . Therefore, Assumption 4 implies that  $c(x, \xi)$  has a continuous density in  $(-\delta, +\delta)$  for any  $x \in \mathcal{O}$ .

**ASSUMPTION 5.** *Let  $\Omega_0 = \{x \in X: p(x) \leq \alpha\}$  and  $\Omega_0^l = \{x \in X: p(x) < \alpha\}$ . Then  $\Omega_0 = cl \Omega_0^l$ .*

Note that  $\Omega_0$  is the set of feasible solutions to Problem (P). Assumption 5 is an assumption on the constraint qualification of Problem (P). When  $\Omega_0$  is a convex set, it is implied by the widely used Slater's condition (e.g., Boyd and Vandenberghe 2004). Assumption 5 is a commonly used condition in nonlinear programming, especially for numerical methods that approximate the optimal solutions by sequences of points in  $\Omega^l$ , e.g., the barrier function method (see, for instance, Zangwill 1969 and Bazaraa et al. 1993). We need this assumption because the method we propose in §3 is also such a numerical method. More discussions on Assumption 5 along with a counterexample are presented in the electronic companion to this paper.

Then, we have the following lemmas that are used repeatedly in the rest of the paper.

**LEMMA 1.** *Suppose that Assumption 4 is satisfied. For any  $x \in X$ ,  $\tilde{p}(x, t)$  is nondecreasing in  $t$  when  $t > 0$ .*

**PROOF.** For any  $t > 0$  and any  $z \in \mathfrak{R}$ ,

$$\begin{aligned} \pi(z, t) &= \psi(z, t) - \phi(z, t) \\ &= \left[1 + \frac{1}{t}z\right] \cdot 1_{(-t, 0]}(z) + 1_{(0, +\infty)}(z). \end{aligned} \quad (5)$$

For any  $t_1 > t_2 > 0$  and any  $z \in \mathfrak{R}$ ,

$$\begin{aligned} \pi(z, t_1) - \pi(z, t_2) &= \left[1 + \frac{1}{t_1}z\right] \cdot 1_{(-t_1, -t_2]}(z) \\ &\quad + \left[\frac{1}{t_1} - \frac{1}{t_2}\right]z \cdot 1_{(-t_2, 0]}(z) \geq 0. \end{aligned}$$

Therefore,  $\pi(z, t)$  is nondecreasing in  $t$  when  $t > 0$ . Because  $\tilde{p}(x, t) = E[\pi(c(x, \xi), t)]$ ,  $\tilde{p}(x, t)$  is also nondecreasing in  $t$  when  $t > 0$ .  $\square$

**LEMMA 2.** *Suppose that Assumptions 1 to 4 are satisfied. Then,  $g_1(x, t)$  is differentiable in  $\mathcal{O} \times (-\delta, \delta)$ , and*

$$\begin{aligned} \nabla_x g_1(x, t) &= E[\nabla_x c_{i^*}(x, \xi) \cdot 1_{(-t, +\infty)}(c(x, \xi))], \\ \frac{\partial}{\partial t} g_1(x, t) &= 1 - F(-t, x), \end{aligned}$$

where  $i^* = \arg \max_{i=1, \dots, m} \{c_i(x, \xi)\}$ .

**PROOF.** By Assumptions 1 to 3,

$$c(x, \xi) = \max_{i=1, \dots, m} c_i(x, \xi)$$

is differentiable with respect to  $x$  w.p.1 and  $\nabla_x c(x, \xi) = \nabla_x c_{i^*}(x, \xi)$  w.p.1 when  $x \in \mathcal{O}$ . Note that  $g_1(x, t) = E\{[t + c(x, \xi)]^+\}$ . Because  $f(x) = (t + x)^+$  is differentiable except

at  $x = -t$  and  $f'(x) = 1_{(-t, +\infty)}(x)$  when  $x \neq -t$ , and  $\Pr\{c(x, \xi) = t\} = 0$  when  $t \in (-\delta, +\delta)$  by Assumption 4, then by Proposition 1 of Broadie and Glasserman (1996),  $g_1(x, t)$  is differentiable in  $\mathcal{O} \times (-\delta, +\delta)$ , and

$$\begin{aligned} \nabla_x g_1(x, t) &= E[\nabla_x c_{i^*}(x, \xi) \cdot 1_{(-t, +\infty)}(c(x, \xi))], \\ \frac{\partial}{\partial t} g_1(x, t) &= E[1_{(-t, +\infty)}(c(x, \xi))] \\ &= \Pr\{c(x, \xi) > -t\} = 1 - F(-t, x). \end{aligned}$$

This concludes the proof of the lemma.  $\square$

REMARK 1. Because  $g_2(x) = g_1(x, 0)$ , Lemma 2 also implies that  $g_2(x)$  is differentiable and

$$\nabla g_2(x) = E[\nabla_x c_{i^*}(x, \xi) \cdot 1_{(0, +\infty)}(c(x, \xi))].$$

Then, we have the following theorem on the equivalence of Problem (DC) and Problem (P).

**THEOREM 1.** *Suppose that Assumptions 1 to 4 are satisfied. Then, Problem (DC) is equivalent to Problem (P).*

PROOF. By Lemma 1,

$$\tilde{p}(x) = \inf_{t>0} \tilde{p}(x, t) = \lim_{t \searrow 0} \tilde{p}(x, t),$$

where  $t \searrow 0$  denotes that  $t$  decreasingly goes to 0. By Lemma 2,

$$\begin{aligned} \lim_{t \searrow 0} \tilde{p}(x, t) &= \lim_{t \searrow 0} \frac{1}{t} [g_1(x, t) - g_1(x, 0)] \\ &= \frac{\partial}{\partial t} g_1(x, 0) = 1 - F(0, x) = p(x). \end{aligned}$$

Then,  $\tilde{p}(x) = p(x)$ . Therefore, Problem (DC) is equivalent to Problem (P). This concludes the proof of the theorem.  $\square$

Theorem 1 is an important result of this paper. It shows that solving Problem (DC) is equivalent to solving Problem (P). In the rest of this paper, we study how to solve Problem (DC).

### 2.4. $\varepsilon$ -Approximation

Note that  $\tilde{p}(x) = \lim_{t \searrow 0} \tilde{p}(x, t)$  by Lemma 1. However,  $\tilde{p}(x, t)$  is not well defined at  $t = 0$ . Therefore, we approximate  $\tilde{p}(x)$  by  $\tilde{p}(x, \varepsilon) = (1/\varepsilon)[g_1(x, \varepsilon) - g_2(x)]$  for a small  $\varepsilon \in (0, \delta)$  where  $\delta$  is defined in Assumption 4, and approximate Problem (DC) by

$$\begin{aligned} (\text{P}_\varepsilon) \quad &\text{minimize } h(x), \\ &\text{subject to } g_1(x, \varepsilon) - g_2(x) \leq \varepsilon\alpha, \\ &x \in X. \end{aligned}$$

By Theorem 1, Problem (DC) is equivalent to Problem (P). Therefore, Problem  $(\text{P}_\varepsilon)$  is also an approximation to Problem (P). In the rest of this subsection, we show that Problem  $(\text{P}_\varepsilon)$  is a good approximation to Problem (P).

Let  $\Omega(\varepsilon) = \{x \in X: g_1(x, \varepsilon) - g_2(x) \leq \varepsilon\alpha\}$  denote the feasible set of Problem  $(\text{P}_\varepsilon)$ . Then, we have the following lemma on the relationship between  $\Omega(\varepsilon)$  and  $\Omega_0$ , which is the feasible set of Problem (P).

**LEMMA 3.** *Suppose that Assumptions 1 to 5 are satisfied. Then,  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) = \Omega_0$ .*

PROOF. By Lemma 1,  $\tilde{p}(x, t)$  is nondecreasing with respect to  $t$ . Then, for any  $\varepsilon_2 \geq \varepsilon_1 > 0$ ,

$$\frac{1}{\varepsilon_1} [g_1(x, \varepsilon_1) - g_2(x)] \leq \frac{1}{\varepsilon_2} [g_1(x, \varepsilon_2) - g_2(x)],$$

which in turn implies that  $\Omega(\varepsilon_2) \subset \Omega(\varepsilon_1)$ . Therefore, it follows from Exercise 4.3 of Rockafellar and Wets (1998) that  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon)$  exists.

We first prove that  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) \subset \Omega_0$ . For any  $x \in \lim_{\varepsilon \searrow 0} \Omega(\varepsilon)$ , there exists  $\varepsilon_k \searrow 0$  and  $x^k \in \Omega(\varepsilon_k)$  such that  $x^k \rightarrow x$ . Because  $x^k \in \Omega(\varepsilon_k)$ , then  $x^k \in X$ , and

$$\frac{1}{\varepsilon_k} [g_1(x^k, \varepsilon_k) - g_2(x^k)] \leq \alpha. \tag{6}$$

By Taylor expansion and Lemma 2, we have

$$\begin{aligned} g_1(x^k, \varepsilon_k) &= g_1(x^k, 0) + \frac{\partial}{\partial t} g_1(x^k, \tilde{\varepsilon}_k) \varepsilon_k \\ &= g_1(x^k, 0) + [1 - F(-\tilde{\varepsilon}_k, x)] \varepsilon_k \end{aligned} \tag{7}$$

for some  $\tilde{\varepsilon}_k \in (0, \varepsilon_k)$ . Combining Equations (6) and (7) and taking  $k \rightarrow +\infty$ , we have  $x \in X$  and  $1 - F(0, x) \leq \alpha$ , which is equivalent to  $p(x) \leq \alpha$ . Therefore,  $x \in \Omega_0$ , which implies that  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) \subset \Omega_0$ .

We then prove that  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) \supset \Omega_0$ . For any  $x \in \Omega'_0$ , because  $p(x) = \lim_{\varepsilon \searrow 0} (1/\varepsilon)[g_1(x, \varepsilon) - g_2(x)]$  and  $p(x) < \alpha$ , then  $(1/\varepsilon)[g_1(x, \varepsilon) - g_2(x)] < \alpha$  for some  $\varepsilon > 0$  small enough. Therefore,  $x \in \Omega(\varepsilon)$ . So we obtain that  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) \supset \Omega'_0$ . Because  $\Omega(\varepsilon)$  is a closed set for any  $\varepsilon > 0$ , then  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon)$  is also a closed set. Then, by Assumption 5,  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) \supset \Omega_0$ .

Therefore,  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) = \Omega_0$ . This concludes the proof of the lemma.  $\square$

For sets  $A, B \subset \mathfrak{R}^d$ , let  $\text{dist}(x, A) = \inf_{x' \in A} \|x - x'\|$  denote the distance from  $x \in \mathfrak{R}^d$  to  $A$ , and

$$\mathbb{D}(A, B) = \sup_{x \in A} \text{dist}(x, B)$$

denote the deviation of the set  $A$  from the set  $B$  (Shapiro et al. 2009). Let  $S(\varepsilon)$  and  $\nu(\varepsilon)$  be the set of optimal solutions and the optimal value of Problem  $(\text{P}_\varepsilon)$ ,  $S_0$  and  $\nu_0$  be the set of optimal solutions and the optimal value of Problem (P). Then, we have the following theorem.

**THEOREM 2.** *Suppose that Assumptions 1 to 5 are satisfied. Then,  $\lim_{\varepsilon \searrow 0} \mathbb{D}(S(\varepsilon), S_0) = 0$  and  $\lim_{\varepsilon \searrow 0} \nu(\varepsilon) = \nu_0$ .*

PROOF. Let  $\bar{h}(x) = h(x) + I_{\Omega_0}(x)$  and  $\bar{h}_\varepsilon(x) = h(x) + I_{\Omega(\varepsilon)}(x)$ , where  $I_A(x) = 0$  if  $x \in A$  and  $I_A(x) = +\infty$  if  $x \notin A$ . By Lemma 3,  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) = \Omega_0$ . Then, by Proposition 7.4(f) of Rockafellar and Wets (1998), we have that  $I_{\Omega(\varepsilon)}(\cdot)$  epi-converges to  $I_{\Omega_0}(\cdot)$  as  $\varepsilon \searrow 0$ . Because  $h(\cdot)$  is continuous, we have that  $\bar{h}_\varepsilon(\cdot)$  epi-converges to  $\bar{h}(\cdot)$  as

$\varepsilon \searrow 0$ . As  $\Omega(\varepsilon)$  and  $\Omega_0$  are compact, we have that  $\bar{h}_\varepsilon(\cdot)$  and  $\bar{h}(\cdot)$  are lower semi-continuous and proper.<sup>1</sup> Then, by Theorem 7.33 of Rockafellar and Wets (1998), we have  $\nu(\varepsilon) \rightarrow \nu_0$  and

$$\limsup_{\varepsilon \searrow 0} S(\varepsilon) \subset S_0. \quad (8)$$

Because  $S(\varepsilon)$  and  $S_0$  are subsets of the compact set  $X$ , they are uniformly compact. By the discussions in Example 4.13 of Rockafellar and Wets (1998), we have that Equation (8) implies  $\lim_{\varepsilon \searrow 0} \mathbb{D}(S(\varepsilon), S_0) = 0$ . This concludes the proof of the theorem.  $\square$

Theorem 2 shows that the optimal solutions of Problem  $(P_\varepsilon)$  provide good approximations to the optimal solutions of Problem  $(P)$  when  $\varepsilon$  is close enough to 0.

However, Problem  $(P_\varepsilon)$  is generally a non-convex problem. Therefore, finding an optimal solution to the problem may be difficult. We often find only KKT points of Problem  $(P_\varepsilon)$  (as the method introduced in §3 does). In the rest of this subsection, we analyze the convergence of the KKT points of Problem  $(P_\varepsilon)$  to the KKT points of Problem  $(P)$  as  $\varepsilon \searrow 0$ .

Let  $\Lambda_0$  and  $\Lambda(\varepsilon)$  denote the sets of KKT pairs of Problems  $(P)$  and  $(P_\varepsilon)$ , respectively, namely

$$\Lambda_0 = \left\{ \begin{array}{l} (x, \lambda) \in \Omega_0 \times \mathfrak{R}_+ : 0 \in \nabla h(x) + \lambda \nabla p(x) + N_X(x), \\ \lambda [p(x) - \alpha] = 0, \end{array} \right\},$$

and

$$\Lambda(\varepsilon) = \left\{ \begin{array}{l} (x, \lambda) \in \Omega(\varepsilon) \times \mathfrak{R}_+ : \\ 0 \in \nabla h(x) + \lambda \left[ \frac{\nabla_x g_1(x, \varepsilon) - \nabla g_2(x)}{\varepsilon} \right] + N_X(x), \\ \lambda \left[ \frac{g_1(x, \varepsilon) - g_2(x)}{\varepsilon} - \alpha \right] = 0, \end{array} \right\},$$

where  $N_X(x)$  denotes the normal cone to  $X$  at  $x$  (Bonnans and Shapiro 2000), and the differentiability of  $p(x)$ ,  $g_1(x, \varepsilon)$  and  $g_2(x)$  is ensured by Assumption 4 and Lemma 2. Then, we have the following theorem that shows the relation between  $\Lambda_0$  and  $\Lambda(\varepsilon)$ .

**THEOREM 3.** *Suppose that Assumptions 1 to 5 are satisfied. Then,  $\limsup_{\varepsilon \searrow 0} \Lambda(\varepsilon) \subset \Lambda_0$ .*

**PROOF.** For any  $(x, \lambda) \in \limsup_{\varepsilon \searrow 0} \Lambda(\varepsilon)$ , there exists  $(x_k, \lambda_k) \in \Lambda(\varepsilon_k)$  such that  $(x_k, \lambda_k) \rightarrow (x, \lambda)$ . The inclusion  $(x_k, \lambda_k) \in \Lambda(\varepsilon_k)$  means

$$0 \in \nabla h(x_k) + \lambda_k \left[ \frac{\nabla_x g_1(x_k, \varepsilon_k) - \nabla g_2(x_k)}{\varepsilon_k} \right] + N_X(x_k), \quad (9)$$

$$\lambda_k \left[ \frac{g_1(x_k, \varepsilon_k) - g_2(x_k)}{\varepsilon_k} - \alpha \right] = 0, \quad \lambda_k \geq 0. \quad (10)$$

By Lemma 2,  $(\partial/\partial t)g_1(x, t) = 1 - F(-t, x)$  when  $x \in X$  and  $t \in (-\delta, \delta)$ . Because  $F(t, x)$  is continuously differentiable by Assumption 4, then  $\nabla_x(\partial/\partial t)g_1(x, t) = -\nabla_x F(-t, x)$  is continuous in  $t$  and  $x$ . Then, by Exercise 6.24 of Marsden and Hoffman (1993),

$$\frac{\partial}{\partial t} \nabla_x g_1(x, t) = \nabla_x \frac{\partial}{\partial t} g_1(x, t) = -\nabla_x F(-t, x).$$

Because  $g_2(x) = g_1(x, 0)$ , by the mean-value theorem, we have

$$\frac{\nabla_x g_1(x_k, \varepsilon_k) - \nabla g_2(x_k)}{\varepsilon_k} = \frac{\partial}{\partial t} \nabla_x g_1(x, \tilde{\varepsilon}_k) = -\nabla_x F(-\tilde{\varepsilon}_k, x)$$

for some  $\tilde{\varepsilon}_k \in (0, \varepsilon_k)$ ,  $k = 1, 2, \dots$ . Because  $F(t, x)$  is continuously differentiable by Assumption 4, then we have that

$$\begin{aligned} \lim_{k \rightarrow +\infty} \frac{\nabla_x g_1(x_k, \varepsilon_k) - \nabla g_2(x_k)}{\varepsilon_k} &= - \lim_{k \rightarrow +\infty} \nabla_x F(-\tilde{\varepsilon}_k, x) \\ &= -\nabla_x F(0, x) = \nabla p(x). \end{aligned} \quad (11)$$

Furthermore, by Lemma 2,

$$\lim_{k \rightarrow +\infty} \frac{g_1(x_k, \varepsilon_k) - g_2(x_k)}{\varepsilon_k} = 1 - F(0, x) = p(x). \quad (12)$$

By Lemma 3, we know that  $\Omega(\varepsilon)$  increases as  $\varepsilon \searrow 0$  and  $\lim_{\varepsilon \searrow 0} \Omega(\varepsilon) = \Omega_0$ . Therefore,  $\Omega(\varepsilon) \subset \Omega_0$  for  $\varepsilon > 0$ . It follows from  $x_k \in \Omega(\varepsilon_k)$  and  $x_k \rightarrow x$  that  $x \in \Omega_0$ . By Proposition 6.6 of Rockafellar and Wets (1998),

$$\limsup_{x_k \rightarrow x} N_X(x_k) = N_X(x) \quad (13)$$

when  $x, x_k \in X$ . Then, by taking  $k \rightarrow +\infty$  in Equations (9) and (10), we obtain that  $(x, \lambda) \in \Lambda_0$ . This concludes the proof of the theorem.  $\square$

To obtain a stronger convergence result of the KKT pairs, we make the following assumption.

**ASSUMPTION 6.** *Suppose that the following regularity condition holds for every feasible point  $x \in \Omega_0$ :*

$$\left. \begin{array}{l} 0 \in \lambda \nabla p(x) + N_X(x) \\ \lambda \geq 0, \lambda [p(x) - \alpha] = 0 \end{array} \right\} \implies \lambda = 0. \quad (14)$$

Assumption 6 is a constraint qualification of the constraints  $x \in X$  and  $p(x) - \alpha \leq 0$ , which ensures the existence of KKT pairs. It is a frequently used condition for a set involving the abstract constraint  $x \in X$ ; see, for instance, the basic constraint qualification defined in Theorem 6.14 of Rockafellar and Wets (1998). Let  $\Sigma$  be the set of all stationary points of Problem  $(P)$ . Then,  $\Sigma \subset X$  is a compact set. For any  $x \in \Sigma$ , it follows from Exercise 6.39 and Example 6.40 of Rockafellar and Wets (1998) that Condition (14) is Robinson constraint qualification, and it is Mangasarian-Fromovitz (MF) constraint qualification when  $X$  is a compact polyhedron.

Then, we have the following theorem that is stronger than Theorem 3. The proof of the theorem is provided in the electronic companion to this paper.

**THEOREM 4.** *Suppose that Assumptions 1 to 6 are satisfied and  $X \subset \mathfrak{R}^d$  is a compact polyhedron. Then,  $\lim_{\varepsilon \searrow 0} \mathbb{D}(\Lambda(\varepsilon), \Lambda_0) = 0$ .*

Theorems 3 and 4 show that the cluster points of the sequence of KKT pairs of Problem  $(P_\varepsilon)$  are KKT pairs of Problem  $(P)$ . Therefore, the KKT points of Problem  $(P_\varepsilon)$  are good approximations to the KKT points of Problem  $(P)$  when  $\varepsilon$  is small. In the rest of this paper, we consider how to find a KKT point of Problem  $(P_\varepsilon)$ .

### 3. Sequential Convex Approximations

Problem  $(P_\varepsilon)$  is a DC program, because the left-hand side of the constraint  $g_1(x, \varepsilon) - g_2(x) - \varepsilon\alpha \leq 0$  is a DC function of  $x$ . To solve the problem, we propose to use a sequence of convex approximations. In §3.1, we first introduce an algorithm to solve this type of DC programs. Then, in §3.2, we show how to apply the algorithm to solve Problem  $(P_\varepsilon)$ . The algorithm starts with an initial feasible solution of Problem  $(P_\varepsilon)$ , we show how to find a good initial solution in §3.3.

#### 3.1. Algorithm SCA

Consider the nonlinear optimization problem of the form

$$\begin{aligned} \text{(DCP)} \quad & \text{minimize } h(x), \\ & \text{subject to } g_1(x) - g_2(x) \leq 0, \\ & x \in X, \end{aligned}$$

where  $X \subset \mathfrak{R}^d$  is a nonempty convex compact set,  $h: \mathfrak{R}^d \rightarrow \mathfrak{R}$ ,  $g_i: \mathfrak{R}^d \rightarrow \mathfrak{R}$ ,  $i = 1, 2$ , are real-valued continuously differentiable and convex functions in a bounded open set  $\mathcal{O} \supset X$ . Note that Problem  $(P_\varepsilon)$  is an example of Problem (DCP).

Let  $\Omega = \{x \in X: g_1(x) - g_2(x) \leq 0\}$  and

$$\Omega_y = \{x \in X: g_1(x) - [g_2(y) + \nabla g_2(y)^T(x - y)] \leq 0\}$$

for any  $y \in X$ . Note that  $g_2(y) + \nabla g_2(y)^T(x - y)$  defines a tangent plane of  $g_2(x)$  at  $x = y$ . Because  $g_2(x)$  is convex, we have for any  $y \in X$ ,

$$g_2(x) \geq g_2(y) + \nabla g_2(y)^T(x - y), \quad \forall x \in X,$$

which implies

$$g_1(x) - g_2(x) \leq g_1(x) - [g_2(y) + \nabla g_2(y)^T(x - y)]. \quad (15)$$

Then,  $\Omega_y \subset \Omega$  for any  $y \in X$ . Furthermore, since  $g_1(x) - [g_2(y) + \nabla g_2(y)^T(x - y)]$  is a convex function of  $x$ , then  $\Omega_y$  is a convex subset of  $\Omega$  for any  $y \in X$ . Define Problem  $(CP(y))$  as

$$(CP(y)) \quad \text{minimize } \{h(x): x \in \Omega_y\}.$$

Then,  $CP(y)$  is a convex conservative approximation of Problem (DCP) for any  $y \in X$ . We suggest using the following algorithm to solve Problem (DCP).

#### Algorithm SCA

*Step 0.* Give  $x_0 \in \Omega$  and set  $k = 0$ .

*Step 1.* Stop if  $x_k$  satisfies the KKT condition of Problem (DCP).

*Step 2.* Solve  $CP(x_k)$  to obtain its optimal solution  $x_{k+1}$ .

*Step 3.* Set  $k = k + 1$  and go to Step 1.

A similar algorithm was proposed by Smola et al. (2005) as an approach to solving DC programs in the form of Problem (DCP). However, they did not provide rigorous analysis on the convergence of the algorithm, although they claim that the algorithm can find a KKT point of Problem (DCP). In the rest of this subsection, we analyze the properties of the algorithm and prove the claim of Smola et al. (2005) in a rigorous way.

**PROPERTY 1.** *If  $\{x_k\}$  is generated by Algorithm SCA for Problem (DCP) starting from  $x_0 \in \Omega$ , then  $\{x_k\} \subset \Omega$ .*

**PROOF.** Note that  $x_1 \in \Omega_{x_0}$ , and  $\Omega_y \subset \Omega$  for any  $y \in X$  by Equation (15). Then  $x_1 \in \Omega$ . Therefore, by the principle of induction, we have  $\{x_k\} \subset \Omega$ .  $\square$

**PROPERTY 2.** *If  $\{x_k\}$  is generated by Algorithm SCA for Problem (DCP) starting from  $x_0 \in \Omega$ , then  $\{h(x_k)\}$  is a convergent nonincreasing sequence.*

**PROOF.** Because  $\Omega_{x_k}$  is a convex compact set and  $h$  is convex,  $CP(x_k)$  has a nonempty compact solution set and  $x_{k+1} \in \arg \min\{h(x): x \in \Omega_{x_k}\}$ . Noting that  $x_k \in \Omega_{x_k}$  for every  $k \geq 1$ , we have that  $h(x_{k+1}) \leq h(x_k)$ . As  $h(\cdot)$  is continuous and  $X$  is compact, we obtain that  $\inf_k \{h(x_k)\}$  is finite, which is greater than or equal to  $\inf_{x \in X} h(x)$ , and  $\lim_{k \rightarrow +\infty} h(x_k) = \inf_k \{h(x_k)\}$ .  $\square$

The next property states that the cluster points of  $\{x_k\}$  are all KKT points of Problem (DCP). To prove it, we need a constraint qualification. We say that Slater's condition holds at  $y \in \Omega$  if  $\text{int}\Omega_y \neq \emptyset$ . Note that Slater's condition is the most commonly used constraint qualification in convex optimization (Boyd and Vandenberghe 2004). The proof of the property is quite lengthy, so we include it in the electronic companion to this paper.

**PROPERTY 3.** *Let  $\{x_k\}$  be the sequence of solutions generated by Algorithm SCA for Problem (DCP) starting from  $x_0 \in \Omega$ . Suppose that  $\bar{x}$  is a cluster point of  $\{x_k\}$  satisfying Slater's condition. Then,  $\bar{x}$  is a KKT point of Problem (DCP). Moreover, if  $h$  is strictly convex in  $\mathcal{O}$ , then  $\{x_k\}$  converges to a KKT point of Problem (DCP).*

By Properties 1 to 3, we see that Algorithm SCA has many desired properties. In the next subsection, we show how to apply it to solve Problem  $(P_\varepsilon)$ .

#### 3.2. Algorithm SCA for Problem $(P_\varepsilon)$

Note that Problem  $(P_\varepsilon)$  is exactly in the form of Problem (DCP), where we need only to define  $g_1(x) = g_1(x, \varepsilon) - \varepsilon\alpha$ . Then we can apply Algorithm SCA to solve Problem  $(P_\varepsilon)$  directly.



By Assumption 1 and the conclusions of Lemma 2, we can verify that Problem  $(P_\varepsilon)$  satisfies the definition of Problem (DCP) when  $\varepsilon$  is small enough. Let  $\{x_k\}$  be the sequence of solutions generated by Algorithm SCA for Problem  $(P_\varepsilon)$  starting from  $x_0 \in \Omega(\varepsilon)$ . By Properties 1 and 2, we have that  $\{x_k\} \subset \Omega(\varepsilon)$  and  $\{h(x_k)\}$  is a convergent nonincreasing sequence when  $\varepsilon$  is small enough.

To apply Property 3 to Problem  $(P_\varepsilon)$ , however, we need to prove that all cluster points of  $\{x_k\}$  satisfy Slater's condition. Let  $\bar{x}$  denote a cluster point of  $\{x_k\}$ . Because  $g_1(x, \varepsilon)$  and  $g_2(x)$  are both continuous in  $x$ , then  $\Omega(\varepsilon)$  is a closed set. Because  $\{x_k\} \subset \Omega(\varepsilon)$ , we have  $\bar{x} \in \Omega(\varepsilon)$ . Then, we need only to prove that  $\text{int } \Omega(\varepsilon)_{\bar{x}} \neq \emptyset$ , which is implied by the conclusion of the following lemma.

**LEMMA 4.** *Suppose that Assumptions 1 to 6 are satisfied. Then,  $\text{int } \Omega(\varepsilon)_y \neq \emptyset$  for any  $y \in \Omega(\varepsilon)$  when  $\varepsilon > 0$  is small enough.*

**PROOF.** By contradiction. Suppose that there exist  $\varepsilon_i \searrow 0$  and  $y_i \in \Omega(\varepsilon_i)$  such that  $\text{int } \Omega(\varepsilon_i)_{y_i} = \emptyset$ . Note that

$$\Omega(\varepsilon)_{y_i} = \{x \in X: g_1(x, \varepsilon_i) - [g_2(y_i) + \nabla g_2(y_i)(x - y_i)] - \varepsilon_i \alpha \leq 0\}.$$

Then, the equality  $\text{int } \Omega(\varepsilon_i)_{y_i} = \emptyset$  implies  $g_1(y_i, \varepsilon_i) - g_2(y_i) - \varepsilon_i \alpha = 0$  and

$$y_i \in \arg \min_{x \in X} \{g_1(x, \varepsilon_i) - [g_2(y_i) + \nabla g_2(y_i)(x - y_i)] - \varepsilon_i \alpha\}. \quad (16)$$

As  $X$  is a convex compact set, the optimization problem of Equation (16) is a convex problem. Then, we have from the necessary optimality condition that

$$-[\nabla_x g_1(y_i, \varepsilon_i) - \nabla g_2(y_i)] \in N_X(y_i),$$

$$g_1(y_i, \varepsilon_i) - g_2(y_i) - \varepsilon_i \alpha = 0,$$

or equivalently

$$\begin{aligned} & -\frac{\nabla_x g_1(y_i, \varepsilon_i) - \nabla_x g_1(y_i, 0)}{\varepsilon_i} \in N_X(y_i), \\ & \frac{g_1(y_i, \varepsilon_i) - g_1(y_i, 0)}{\varepsilon_i} - \alpha = 0. \end{aligned} \quad (17)$$

Because  $\{y_i\} \subset X$  and  $X$  is compact,  $\{y_i\}$  has a cluster point, say  $\bar{y} \in \Omega_0$ . Assume that there is a subsequence  $\{y_{k_j}\}$  such that  $y_{k_j} \rightarrow \bar{y}$ . Letting  $j \rightarrow +\infty$  and by Equations (11) to (13), we have

$$-\nabla p(\bar{y}) \in N_X(\bar{y}), \quad p(\bar{y}) - \alpha = 0, \quad \bar{y} \in \Omega_0,$$

which implies that  $\lambda = 1$  is a solution of

$$0 \in \lambda \nabla p(x) + N_X(x), \quad \lambda \geq 0, \quad \lambda[p(x) - \alpha] = 0$$

when  $x = \bar{y} \in \Omega_0$ . Then, it contradicts Assumption 6. This concludes the proof of the lemma.  $\square$

Note that Lemma 4 shows that  $\text{int } \Omega(\varepsilon)_y \neq \emptyset$  for any  $y \in \Omega(\varepsilon)$ . Then, it also holds for  $y = \bar{x}$ . Therefore, the conclusions of Property 3 also hold when Algorithm SCA is applied to solve Problem  $(P_\varepsilon)$  when  $\varepsilon$  is small enough.

For completeness, we summarize the three properties for Problem  $(P_\varepsilon)$  in the following theorem.

**THEOREM 5.** *Let  $\{x_k\}$  be the sequence of solutions generated by Algorithm SCA for Problem  $(P_\varepsilon)$  starting from  $x_0 \in \Omega(\varepsilon)$ . Suppose that Assumptions 1 to 6 are satisfied. Then, for any  $\varepsilon > 0$  small enough,  $\{x_k\} \subset \Omega(\varepsilon)$ ,  $\{h(x_k)\}$  is a convergent nonincreasing sequence, all cluster points of  $\{x_k\}$  are KKT points of Problem  $(P_\varepsilon)$ . Furthermore,  $\{x_k\}$  converges to a KKT point of Problem  $(P_\varepsilon)$  if  $h$  is strictly convex.*

### 3.3. Initial Solutions for Problem $(P_\varepsilon)$

To apply Algorithm SCA, we need an initial solution  $x_0 \in \Omega(\varepsilon)$ . In this subsection, we provide two natural choices.

In the first choice, we let  $\Omega_0(\varepsilon) = \{x \in X: g_1(x, \varepsilon) \leq \varepsilon \alpha\}$ . Note that  $g_2(x) = \mathbb{E}[c(x, \xi)]^+ \geq 0$  for all  $x \in X$ . Then,  $\Omega_0(\varepsilon) \subset \Omega(\varepsilon)$ . Furthermore,  $\Omega_0(\varepsilon)$  is a convex set since  $g_1(x, \varepsilon)$  is a convex function of  $x$  when  $x \in \mathcal{C}$ . Then,

$$\begin{aligned} & \text{minimize } h(x), \\ & \text{subject to } x \in \Omega_0(\varepsilon) \end{aligned}$$

is a convex optimization problem. Let  $x_\varepsilon \in \arg \min\{h(x): x \in \Omega_0(\varepsilon)\}$ , we have  $x_\varepsilon \in \Omega(\varepsilon)$ .

In the second choice, we let  $\Omega_{\text{CVaR}} = \{x \in X: \text{CVaR}_{1-\alpha}(c(x, \xi)) \leq 0\}$ . By the discussions in §2.1,

$$\Omega_{\text{CVaR}} = \left\{x \in X: \inf_{t>0} \frac{1}{t} g_1(x, t) \leq \alpha, x \in X\right\}.$$

Let  $x_{\text{CVaR}} \in \arg \min\{h(x): x \in \Omega_{\text{CVaR}}\}$ , which is the optimal solution of the CVaR approximation of Rockafellar and Uryasev (2000). Let  $\varepsilon^* = q_{1-\alpha}(c(x_{\text{CVaR}}, \xi))$ , which is the  $1 - \alpha$  quantile of  $c(x_{\text{CVaR}}, \xi)$ . By Pflug (2000),  $\varepsilon^* > 0$  and

$$\inf_{t>0} \frac{1}{t} g_1(x_{\text{CVaR}}, t) = \frac{1}{\varepsilon^*} g_1(x_{\text{CVaR}}, \varepsilon^*).$$

Then,  $x_{\text{CVaR}} \in \{x \in X: g_1(x, \varepsilon^*) \leq \varepsilon^* \alpha\}$ . Since  $g_2(x) \geq 0$ , then

$$x_{\text{CVaR}} \in \{x \in X: g_1(x, \varepsilon^*) - g_2(x) \leq \varepsilon^* \alpha\} = \Omega(\varepsilon^*).$$

By Lemma 1,  $\Omega(\varepsilon^*) \subset \Omega(\varepsilon)$  for any  $0 < \varepsilon \leq \varepsilon^*$ . Then,  $x_{\text{CVaR}} \in \Omega(\varepsilon)$  for any  $0 < \varepsilon \leq \varepsilon^*$ . Therefore, one may first solve the CVaR approximation and find  $x_{\text{CVaR}}$  and then select  $\varepsilon \in (0, \varepsilon^*]$ . Then,  $x_{\text{CVaR}}$  may be used as an initial solution to Algorithm SCA.

When we let  $x_0 = x_{\text{CVaR}}$ , the sequence of solutions  $\{x_k\}$  generated by Algorithm SCA are improving and at least as good as the CVaR approximation, which is the “best”

convex conservative approximation. Furthermore,  $\{x_k\}$  converges to the set of KKT points of Problem (P<sub>ε</sub>) by Theorem 5, which converges to the set of KKT points of Problem (P) as  $\varepsilon \searrow 0$  by Theorem 4. Compared to other approximation algorithms, e.g., CVaR approximation, quadratic approximation, Bernstein approximation and scenario analysis, which only find a (good) feasible solution to Problem (P), our algorithm has more desirable properties. If Problem (P) is convex (even though it may not be verifiable), our algorithm converges to its global optimal solution, while others do not.

### 4. A Gradient-Based Monte Carlo Method

To implement Algorithm SCA, we need to repeatedly solve minimize  $h(x)$ ,

$$\text{subject to } g_1(x, \varepsilon) - [g_2(y) + \nabla g_2(y)^T(x - y)] \leq \varepsilon \alpha, \quad (18)$$

$$x \in X$$

for different  $y$  values. Though Problem (18) is a convex optimization problem, it is difficult to solve because we generally do not have the closed-form expressions of  $g_1(x, \varepsilon)$ ,  $g_2(x)$ , and  $\nabla g_2(y)$ . To overcome this difficulty, we propose to use a Monte Carlo method.

For simplicity of the notation, we let

$$g(x) = g_1(x, \varepsilon) - [g_2(y) + \nabla g_2(y)^T(x - y)].$$

Then, by the definitions of  $g_1$ ,  $g_2$  and Lemma 2, we have

$$g(x) = E\{[c(x, \xi) + \varepsilon]^+ - [E\{[c(y, \xi)]^+ + E\{[\nabla_x c_{i^*}(y, \xi) \cdot 1_{(0, +\infty)}(c(y, \xi))\}]^T(x - y)]\}, \quad (19)$$

where

$$c(x, \xi) = \max\{c_i(x, \xi)\} \quad \text{and} \quad i^* = \arg \max_{i=1, \dots, m} \{c_i(y, \xi)\}.$$

Let  $\xi_1, \dots, \xi_n$  denote an independent and identically distributed (i.i.d.) sample of  $\xi$ . Let  $\bar{g}_2(y) = (1/n) \sum_{l=1}^n c(y, \xi_l)$  and  $\bar{\nabla} g_2(y) = (1/n) \sum_{l=1}^n \nabla_x c_{i^*}(y, \xi_l) \cdot 1_{(-t, +\infty)}(c(y, \xi_l))$ . Then, a natural estimator of  $g(x)$  is

$$\bar{g}(x) = \frac{1}{n} \sum_{l=1}^n [c(x, \xi_l) + \varepsilon]^+ - [\bar{g}_2(y) + \bar{\nabla} g_2(y)^T(x - y)]. \quad (20)$$

We suggest solving

$$\text{(MC) minimize } h(x),$$

$$\text{subject to } \bar{g}(x) \leq \varepsilon \alpha, \quad x \in X,$$

and using its optimal solution to approximate the optimal solution of Problem (18). Let  $S$  and  $\nu^*$  denote the set of optimal solutions and the optimal objective value of Problem (18), and  $\hat{S}_n$  and  $\hat{\nu}_n^*$  denote the set of optimal solutions and the optimal objective value of Problem (MC). Then there are two critical issues when we use this approximation. First, do  $\hat{S}_n$  and  $\hat{\nu}_n^*$  converge to  $S$  and  $\nu^*$ ? Second, how do we solve Problem (MC) efficiently?

To answer the first question, we have the following theorem.

**THEOREM 6.** *Suppose that Assumptions 1 to 6 are satisfied. When  $\varepsilon > 0$  is small enough,  $\mathbb{D}(\hat{S}_n, S) \rightarrow 0$  w.p.1 and  $\hat{\nu}_n^* \rightarrow \nu^*$  w.p.1 as  $n \rightarrow \infty$ .*

**PROOF.** Note that we may write  $g(x)$  and  $\bar{g}(x)$  as

$$g(x) = E\{[c(x, \xi) + \varepsilon]^+ - [c(y, \xi)]^+ - [\nabla_x c_{i^*}(y, \xi) \cdot 1_{(0, +\infty)}(c(y, \xi))]^T(x - y)\}, \quad (21)$$

and

$$\bar{g}(x) = \frac{1}{n} \sum_{l=1}^n \{[c(x, \xi_l) + \varepsilon]^+ - [c(y, \xi_l)]^+ - [\nabla_x c_{i^*}(y, \xi_l) \cdot 1_{(0, +\infty)}(c(y, \xi_l))]^T(x - y)\}.$$

Then,  $\bar{g}(x)$  is the sample average approximation of  $g(x)$ . By the strong law of large numbers (Durrett 2005),  $\bar{g}(x) \rightarrow g(x)$  w.p.1 for any fixed  $x \in X$ . Furthermore, because the integrand of Equation (21) is a convex function of  $x$  for any  $\xi \in \Xi$  when  $x \in \mathcal{C}$ , by Theorem 7.50 of Shapiro et al. (2009),  $\bar{g}(x)$  converges to  $g(x)$  uniformly on  $X$  w.p.1 as  $n \rightarrow \infty$ , i.e.,

$$\sup_{x \in X} |\bar{g}(x) - g(x)| \rightarrow 0 \quad \text{w.p.1 as } n \rightarrow \infty.$$

Also, by Lemma 4, Slater’s condition holds for Problem (18) when Assumptions 1 to 6 are satisfied and  $\varepsilon > 0$  is small enough. Then, by Theorem 5.5 of Shapiro et al. (2009) and the discussions followed the theorem, the conclusions of our theorem hold.  $\square$

To answer the second question, we propose two methods to solve Problem (MC). In the first method, based on Equation (20) and the definition of  $c(x, \xi)$ , we reformulate Problem (MC) as

$$\text{minimize } h(x),$$

$$\text{subject to } c_i(x, \xi_l) \leq z_l, \quad i = 1, \dots, m, l = 1, \dots, n,$$

$$\frac{1}{n} \sum_{l=1}^n z_l - [\bar{g}_2(y) + \bar{\nabla} g_2(y)^T(x - y)] \leq \varepsilon \alpha,$$

$$z_l \geq 0, \quad l = 1, \dots, n,$$

$$x \in X. \quad (22)$$

Note that Problem (22) is similar to the formulation of the sample CVaR problem of Rockafellar and Uryasev (2000). It is a convex optimization problem. Furthermore, it is a linear program if  $c_i(x, \xi)$  are linear functions of  $x$  for all  $i = 1, \dots, m$ . However, Problem (22) is often slow to solve because of the large numbers of decision variables and constraints, especially when the sample size  $n$  is large.

To efficiently solve Problem (MC) when  $n$  is large, we propose a second method. By Equation (19) and Lemma 2,

$$\nabla g(x) = E[\nabla_x c_{j^*}(x, \xi) \cdot 1_{(-\varepsilon, +\infty)}(c(x, \xi))] - E[\nabla_x c_{i^*}(y, \xi) \cdot 1_{(0, +\infty)}(c(y, \xi))],$$

where

$$j^* = \arg \max_{j=1, \dots, m} \{c_j(x, \xi)\} \quad \text{and} \quad i^* = \arg \max_{i=1, \dots, m} \{c_i(y, \xi)\}.$$

Then,  $\nabla g(x)$  can be estimated by

$$\bar{\nabla} g(x) = \frac{1}{n} \sum_{l=1}^n [\nabla_x c_{j^*}(x, \xi_l) \cdot 1_{(-\varepsilon, +\infty)}(c(x, \xi_l)) - \nabla_x c_{i^*}(y, \xi_l) \cdot 1_{(0, +\infty)}(c(y, \xi_l))].$$

Although  $g(x)$  is differentiable in  $\mathcal{O}$ ,  $\bar{g}(x)$  is not when  $\xi_1, \dots, \xi_n$  are given. It is only piecewise differentiable. At the points where  $\bar{g}(x)$  is differentiable,  $\nabla \bar{g}(x) = \bar{\nabla} g(x)$ ; at the points where  $\bar{g}(x)$  is not differentiable,  $\bar{\nabla} g(x)$  is a subgradient of  $\bar{g}(x)$ . Therefore, one may use a subgradient-based algorithm (see, for instance, Freund 2004) to solve Problem (MC). In this paper, however, we suggest using an approximation method. Because  $\bar{g}(x)$  converges to  $g(x)$ , which is continuously differentiable as  $n \rightarrow \infty$ , we may approximate  $\bar{\nabla} g(x)$  as a smooth function when  $n$  is large and may use  $\bar{\nabla} g(x)$  as its gradient. Then, we can use gradient-based algorithms to solve Problem (MC) directly. Note that this method can also be viewed as directly solving Problem (18) with estimated  $g(x)$  and  $\nabla g(x)$ .

When we use the gradient-based method to solve Problem (MC), the samples are used only to compute  $\bar{g}(x)$  and  $\bar{\nabla} g(x)$ , which is an  $O(n)$  operation. The method is generally much faster than the first method that solves Problem (22). Hong and Liu (2009) compared the two methods for the CVaR approximation problem through numerical examples. They reported that the two methods find solutions with similar quality, but the gradient-based method is at least an order-of-magnitude faster when  $n$  is of moderate or large size, e.g.,  $n \geq 2,000$ . In our numerical experiments, we observe good performances of the gradient-based method as well.

## 5. Numerical Illustration

In this section, we consider two JCCP problems, a norm optimization problem and a network optimization problem. We use them to illustrate the performances of our method, and we compare our method to the CVaR approximation and the scenario approach.

### 5.1. A Norm Optimization Problem

Let  $x = (x_1, \dots, x_d)^T$  denote a  $d$ -dimensional vector in  $\mathfrak{R}^d$ , and let  $\xi = (\xi_1, \dots, \xi_m)$ , with  $\xi_i = (\xi_{i1}, \dots, \xi_{id})^T$  for any  $i = 1, \dots, m$ , be a  $d \times m$  matrix of random variables. Let  $\|x\|_1$  and  $\|x\|$  denote the 1-norm and 2-norm of  $x$ , respectively, i.e.,  $\|x\|_1 = \sum_{j=1}^d |x_j|$  and  $\|x\| = (\sum_{j=1}^d x_j^2)^{1/2}$ , and let  $\xi_i \circ x = (\xi_{i1}x_1, \dots, \xi_{id}x_d)^T$  denote the Hadamard product (or entrywise product) of  $\xi_i$  and  $x$ . We are interested in solving the following problem:

$$\begin{aligned} & \text{maximize} && \|x\|_1, \\ & \text{subject to} && \Pr\{\|\xi_i \circ x\| \leq 10, i = 1, \dots, m\} \geq 1 - \alpha, \\ & && x_j \geq 0, \quad j = 1, \dots, d. \end{aligned} \quad (23)$$

We may reformulate Problem (23) as

$$\begin{aligned} & \text{minimize} && -\sum_{j=1}^d x_j, \\ & \text{subject to} && \Pr\left\{\sum_{j=1}^d \xi_{ij}^2 x_j^2 \leq 100, i = 1, \dots, m\right\} \geq 1 - \alpha, \\ & && x_j \geq 0, \quad j = 1, \dots, d. \end{aligned} \quad (24)$$

Note that Problem (24) is a JCCP as defined in Problem (P).

Let  $c_i(x, \xi) = \sum_{j=1}^d \xi_{ij}^2 x_j^2 - 100$  for all  $i = 1, \dots, m$ . For any  $x \neq 0$ ,  $c_i(x, \xi)$  is a continuous random variable and  $c_i(x, \xi) = c_j(x, \xi)$  with probability 0. Therefore, Assumption 3 can be satisfied easily. When  $x = 0$ ,  $c_i(x, \xi) = -100$  for all  $i = 1, \dots, m$ . By the definition of differentiability, for any  $j = 1, \dots, d$ ,

$$\begin{aligned} \frac{\partial}{\partial x_j} c(0, \xi) &= \lim_{\delta \rightarrow 0} \frac{1}{\delta} [c(e_j \delta, \xi) - c(0, \xi)] \\ &= \lim_{\delta \rightarrow 0} \frac{1}{\delta} \max_{i=1, \dots, m} (\xi_{ij}^2 \delta^2) = \lim_{\delta \rightarrow 0} \max_{i=1, \dots, m} \xi_{ij}^2 \cdot \delta = 0, \end{aligned}$$

where  $e_j$  denotes the  $i$ th column of a  $d \times d$  identity matrix. Therefore,  $c(x, \xi)$  is differentiable at  $x = 0$  for any  $\xi$ , and Assumption 3 is satisfied.

In the rest of this subsection, we apply Algorithm SCA with the gradient-based Monte Carlo method to solve this problem.

**5.1.1. Independent Case.** We consider the case where  $\xi_{ij}$ ,  $i = 1, \dots, m$  and  $j = 1, \dots, d$ , are independent and identically distributed standard normal random variables. We will call this case *independent case* in the rest of this section. Note that convexity of Problem (23) in this situation is not clear. However, by symmetry, the optimal solution satisfies  $x_1 = \dots = x_d$ . Then,

$$\Pr\left\{\sum_{j=1}^d \xi_{ij}^2 x_j^2 \leq 100, i = 1, \dots, m\right\} = \left[\Pr\left\{x_1^2 \sum_{j=1}^d \xi_{1j}^2 \leq 100\right\}\right]^m.$$

Note that  $\sum_{j=1}^d \xi_{1j}^2$  follows a chi-square distribution with  $d$  degrees of freedom. Let  $F_{\chi_d^2}(\cdot)$  denote its distribution function. Then, the joint chance constraint of Problem (24) is equivalent to

$$F_{\chi_d^2}\left(\frac{100}{x_1^2}\right) \geq (1 - \alpha)^{1/m}.$$

Let  $1 - \beta = (1 - \alpha)^{1/m}$  and let  $F_{\chi_d^2}^{-1}(\cdot)$  denote the inverse distribution function of a chi-square distribution with  $d$  degrees of freedom. Then, it is clear that the optimal solution  $x^*$  of Problem (24) is

$$x_1^* = \dots = x_d^* = \frac{10}{\sqrt{F_{\chi_d^2}^{-1}(1 - \beta)}}.$$

We apply our gradient-based Monte Carlo method directly to solve Problem (24) without exploring its special structure (e.g., the independence of  $\xi$ ), and we use  $x^*$  as a benchmark to evaluate the performances of our method.


We set  $d = 10$ ,  $m = 10$ , and  $\alpha = 0.1$ . Then, the optimal solution of Problem (24) is  $x_1^* = \dots = x_d^* = 2.08$  and the optimal objective value is  $f^* = -20.82$ . We set  $\varepsilon = 0.05^2$  and use a sample size  $n = 10,000$ . We use both  $x_{\text{CVaR}}$  and  $x_\varepsilon$  as initial solutions to compare their performances, and we stop the algorithm if the difference between the objective values of two consecutive iterations is less than or equal to 0.01. We implement the algorithm in Matlab and use Matlab's own nonlinear optimization solver `fmincon` to solve the optimization problem in each iteration with the estimated constraint values and estimated gradients. The programs were run on a desktop computer with Intel Duo Core CPU (3.16 GHz, 3.16 GHz) and 4 GB of RAM.

Although our algorithm uses Monte Carlo samples, we find the performances of the algorithm are very stable. We run the algorithm 100 replications, it always converges to similar solutions. We report the typical performances of the algorithm in Figure 3. In the left panel of Figure 3, we plot the objective values of all iterations. From the plot, we can see that the algorithm converges to the optimal objective value from both  $x_{\text{CVaR}}$  and  $x_\varepsilon$ , even though the convexity of the problem is not clear. Furthermore, we can see that  $x_{\text{CVaR}}$


is a better initial solution, because the algorithm converges faster from it than from  $x_\varepsilon$ . In the right panel of Figure 3, we plot the values of left-hand side of the joint chance constraint, estimated using the sample, for all iterations. From the plot, we see that the joint chance constraint becomes tight when our algorithm stops, while it is not tight at  $x_\varepsilon$  and  $x_{\text{CVaR}}$ . The algorithm typically requires about 20 iterations to converge when  $x_{\text{CVaR}}$  is used as the initial solution and about 35 iterations when  $x_\varepsilon$  is used. The CPU time of each iteration is on average 6.8 seconds with a range of 4.4 to 12.2 seconds.

**5.1.2. Dependent Case.** We also considered the case where  $\xi_{ij}$ ,  $i = 1, \dots, m$  and  $j = 1, \dots, d$ , are dependent. We let  $\xi_{ij}$  be a normal random variable with mean  $j/d$  and variance 1,  $\text{Cov}(\xi_{ij}, \xi_{i'j}) = 0.5$  when  $i \neq i'$  and  $\text{Cov}(\xi_{ij}, \xi_{i'j'}) = 0$  when  $j \neq j'$ . We will call this case *dependent case* in the rest of this subsection. The optimal solution of this case is no longer known. However, we can still apply our method to solve the problem. With the same setting of the parameters as in the independent case, we report the performances of the algorithm in Figure 4. From the

**Figure 3.** Performance of algorithm SCA for independent case.



**Figure 4.** Performance of Algorithm SCA for dependent case



plots, we can see that changing the dependence structure of  $\xi$  does not alter the performances of our algorithm. Both initial solutions lead the algorithm to converge to the same objective value, and the joint chance constraint becomes tight when the algorithm stops. The number of iterations and the CPU time in this case are also similar to the ones in the independent case.

**5.1.3. Scenario Approach.** We also implement the scenario approach for this problem with  $d = m = 10$  and  $\alpha = 0.1$ . By Calafiore and Campi (2006), we let the sample size equal 1,038. Then, by the formulation of Problem (2), we need to solve an optimization problem with 10,380 constraints. We also use Matlab's nonlinear optimization solver `fmincon` to solve the problem. We run the algorithm 100 replications. To our surprise, the problem can be solved very efficiently (0.74 seconds on average) even though there are more than 10,000 constraints. This is because the number of active constraints is only about 10 for all the replications that we have tried. However, the solutions are not stable, i.e., they are drastically different from replication to replication due to the randomness in the sample. For the independent case, the average objective value is  $-17.6$ , with the best being  $-18.6$ , and the estimated left-hand side value of the joint chance constraint is in the range of 0.982 to 0.999. For the dependent case, the average objective value is  $-15.7$  with the best being  $-16.7$  and the estimated left-hand side value of the joint chance constraint being in the range of 0.984 to 0.999. For both cases, the solutions found by the scenario approach are too conservative, and they are significantly worse than the solutions found by the CVaR approximation and our method, even though it is the fastest among all three.

## 5.2. A Network Optimization Problem

A generalized network flow problem (GNFP) is an extension of the classical network flow problem. In a GNFP, the flow on an arc is subject to change. If we use  $x_{ij}$  to denote the flow on arc  $e_{ij}$  when the flow leaves node  $i$ , then the flow will become  $\eta_{ij}x_{ij}$  when it arrives at node  $j$ , where  $0 \leq \eta_{ij} \leq 1$  is a parameter indicating the changing rate. The GNFP has many useful applications. For example, in telecommunication, it models the packet loss on an unreliability transmission link; in production planning, it models the yield rate when one material is converted to another material. In our computational experiments, we consider a GNFP for electricity distribution where  $\eta_{ij}$  is used to model the power loss incurred on a transmission line (Jensen and Bard 2003).

In the problem, there are three electricity generating stations that serve ten different areas. Each station has a different generating cost and a capacity, and each area has a demand. The electricity transmission from a station to an area or to another station is subject to a random percentage loss, which is denoted by  $\eta_{ij}$ . We model  $\eta_{ij}$  by a beta distribution, whose mean and variance depend on the distance

between  $i$  and  $j$ , and  $0 \leq \eta_{ij} \leq 1$ . The problem is to use the minimum cost to serve all areas where the probability of under-supply for any area or generating station is below a certain bound.

Let  $c_i$ ,  $l_i$ , and  $u_i$  denote the marginal generation cost, the minimum generation amount, and the capacity at station  $i$ , respectively, let  $d_k$  denote the demand at area  $k$ , and let  $x_i$ ,  $y_{ik}$ , and  $z_{ij}$  denote the electricity generated at station  $i$ , the electricity transmitted from station  $i$  to area  $k$ , and the electricity transmitted from station  $i$  to station  $j$ , respectively, for all  $i, j = 1, 2, 3$ ,  $i \neq j$ , and  $k = 1, \dots, 10$ . Note that  $x_i$ ,  $y_{ik}$ , and  $z_{ij}$  are decision variables. Then, the problem can be formulated as a JCCP as follows:

$$\begin{aligned} & \text{minimize} && \sum_{i=1}^3 c_i x_i, \\ & \text{subject to} && \Pr \left\{ x_i + \sum_{j \neq i} \eta_{ji} z_{ji} \geq \sum_{k=1}^{10} y_{ik} + \sum_{j \neq i} z_{ij}, i = 1, 2, 3, \right. \\ & && \left. \sum_i \eta_{ik} y_{ik} \geq d_k, k = 1, \dots, 10, \right\} \geq 1 - \alpha \\ & && l_i \leq x_i \leq u_i, y_{ik} \geq 0, z_{ij} \geq 0, \\ & && i, j = 1, 2, 3, i \neq j, k = 1, \dots, 10. \end{aligned}$$

For the test problem we considered, we set  $\alpha = 0.1$  and use a sample size of 10,000. We compare the CVaR approximation and our method and run both algorithms 100 times. The (estimated) optimal cost from the CVaR approximation is  $2.60 \times 10^6$ , while the (estimated) optimal cost from our method is  $2.07 \times 10^6$ , which is roughly a 20% reduction of from the CVaR solution.

## 6. Conclusion and Future Research

In this paper, we propose a sequential convex approximation algorithm that directly solves the stochastic optimization problems with a joint chance constraint. In each iteration of the algorithm, it solves a convex optimization problem using a Monte Carlo method. We have shown that the algorithm finds a KKT point in the limit if the sample size of the Monte Carlo method goes to infinity and the parameter  $\varepsilon$  goes down to zero. Among the algorithms that have been proposed to solve this type of problem, to the best of our knowledge, our algorithm is the first one that has provable convergence to the set of KKT points. It is conceptually more attractive than conservative convex approximation algorithms and the scenario approach, which find (good) feasible solutions that nevertheless satisfy no optimality conditions.

To apply this algorithm for practical problems, especially large-scale practical problems, there are several impediments. First, the algorithm is generally slower than other convex approximation algorithms because it needs to solve a sequence of convex optimization problems and because it uses a Monte Carlo method in each iteration. How to speed up the algorithm is a very important topic for future

research. Second, our method requires the full joint distribution of the uncertain parameters in order to generate Monte Carlo samples. However, specifying a full joint distribution for a large number of parameters is often very challenging in practice. Different robust optimization algorithms have been proposed to address this issue. A future research topic is how to apply our algorithm under the robust optimization framework to find better solutions.

## 7. Electronic Companion

An electronic companion to this paper is available as part of the online version that can be found at <http://or.journal.informs.org/>.

## Endnotes

1. A function  $f: \mathfrak{N}^d \rightarrow \mathfrak{R} \cup \{\pm\infty\}$  is lower semi-continuous at  $x_0 \in \mathfrak{N}^d$  if  $f(x_0) \leq \liminf_{x \rightarrow x_0} f(x)$ . It is lower semi-continuous if it is lower semi-continuous at every  $x \in \mathfrak{N}^d$ . It is a proper function if  $f(x) > -\infty$  for every  $x \in \mathfrak{N}^d$  and there is at least one point  $x \in \mathfrak{N}^d$  such that  $f(x) < +\infty$ .
2. Based on the convergence analysis, we might want to set  $\varepsilon$  small to reduce the bias. However, extremely small  $\varepsilon$  might cause numerical problems and might require longer time to solve the subproblem in each iteration. Given the error in Monte Carlo estimation, we do not suggest setting  $\varepsilon$  extremely small. Finding the optimal setting of  $\varepsilon$  is an important problem for future research.

## Acknowledgments

The authors thank Area Editor Shane Henderson, the associate editor, and three anonymous referees for their helpful comments and suggestions. This research was partially supported by Hong Kong Research Grants Council grant CERG 613907.

## References

- Bazaraa, M. S., H. D. Sherali, C. M. Shetty. 1993. *Nonlinear Programming: Theory and Algorithms*, 2nd ed. John Wiley & Sons, New York.
- Ben-Tal, A., A. Nemirovski. 2000. Robust solutions of linear programming problems contaminated with uncertain data. *Math. Programming* **88**(3) 411–424.
- Bertsimas, D., M. Sim. 2004. Price of robustness. *Oper. Res.* **52**(1) 35–53.
- Bonnans, J. F., A. Shapiro. 2000. *Perturbation Analysis of Optimization Problems*. Springer, New York.
- Boyd, S., L. Vandenberghe. 2004. *Convex Optimization*. Cambridge University Press, Cambridge, UK.
- Broadie, M., P. Glasserman. 1996. Estimating security price derivatives using simulation. *Management Sci.* **42**(2) 269–285.
- Calafiore, G., M. C. Campi. 2005. Uncertain convex programs: Randomized solutions and confidence levels. *Math. Programming* **102**(1) 25–46.
- Calafiore, G., M. C. Campi. 2006. The scenario approach to robust control design. *IEEE Trans. Automatic Control* **51**(5) 742–753.
- Charnes, A., W. W. Cooper, G. H. Symonds. 1958. Cost horizons and certainty equivalents: An approach to stochastic programming of heating oil. *Management Sci.* **4**(3) 235–263.
- Chen, W., M. Sim, J. Sun, C.-P. Teo. 2010. From CVaR to uncertainty set: Implications in joint chance-constrained optimization. *Oper. Res.* **58**(2) 470–485.
- de Farias, D. P., B. Van Roy. 2004. On constraint sampling in the linear programming approach to approximate dynamic programming. *Math. Oper. Res.* **29**(3) 462–478.
- Dentcheva, D., B. Lai, A. Ruszczyński. 2004. Dual methods for probabilistic optimization problems. *Math. Methods Oper. Res.* **60**(2) 331–346.
- Durrett, R. 2005. *Probability: Theory and Examples*, 3rd ed. Duxbury Press, Belmont, MA.
- Erdogan, E., G. Iyengar. 2006. Ambiguous chance constrained problems and robust optimization. *Math. Programming* **107**(1) 37–61.
- Freund, R. M. 2004. Subgradient optimization, generalized programming, and nonconvex duality. Technical report, Massachusetts Institute of Technology, Cambridge, MA.
- Henrion, R. 2007. Structural properties of linear probabilistic constraints. *Optimization* **56**(4) 425–440.
- Henrion, R., C. Strugarek. 2008. Convexity of chance constraints with independent random variables. *Comput. Optim. Appl.* **41**(2) 263–276.
- Hong, L. J. 2009. Estimating quantile sensitivities. *Oper. Res.* **57**(1) 118–130.
- Hong, L. J., G. Liu. 2009. Simulating sensitivities of conditional value-at-risk. *Management Sci.* **55**(2) 281–293.
- Hong, L. J., G. Liu. 2010. Pathwise estimation of probability sensitivities through terminating or steady-state simulations. *Oper. Res.* **58**(2) 357–370.
- Jensen, P. A., J. F. Bard. 2003. *Operations Research: Models and Methods*. John Wiley & Sons, New York.
- Lagoa, C. M., X. Li, M. Sznajder. 2005. Probabilistically constrained linear programs and risk-adjusted controller design. *SIAM J. Optim.* **15**(3) 938–951.
- Luedtke, J., S. Ahmed. 2008. A sample approximation approach for optimization with probabilistic constraints. *SIAM J. Optim.* **19**(2) 674–699.
- Luedtke, J., S. Ahmed, G. Nemhauser. 2007. An integer programming approach for linear programs with probabilistic constraints. *Lecture Notes Comput. Sci.* **4513** 410–423.
- Marsden, J., M. Hoffman. 1993. *Elementary Classical Analysis*, 2nd ed. Freeman, New York.
- Miller, L. B., H. Wagner. 1965. Chance-constrained programming with joint constraints. *Oper. Res.* **13**(6) 930–945.
- Nemirovski, A., A. Shapiro. 2006. Convex approximations of chance constrained programs. *SIAM J. Optim.* **17**(4) 969–996.
- Pflug, G. 2000. Some remarks on the value-at-risk and the conditional value-at-risk. S. Uryasev, ed. *Probabilistic Constrained Optimization: Methodology and Applications*. Kluwer, Dordrecht, The Netherlands.
- Prékopa, A. 1970. On probabilistic constrained programming. *Proc. Princeton Sympos. Math. Programming*, Princeton University Press, Princeton, NJ, 113–138.
- Prékopa, A. 2003. Probabilistic programming. A. Ruszczyński, A. Shapiro, eds. *Stochastic Programming*, Vol. 10. Handbooks in Operations Research and the Management Sciences. Elsevier, Amsterdam.
- Prékopa, A., T. Rapcsák, I. Zsuffa. 1978. Serially linked reservoir system design using stochastic programming. *Water Resources Res.* **14**(4) 672–678.
- Rockafellar, R. T., S. Uryasev. 2000. Optimization of conditional value-at-risk. *J. Risk* **2**(3) 21–41.
- Rockafellar, R. T., R. J.-B. Wets. 1998. *Variational Analysis*. Springer-Verlag, New York.
- Shapiro, A., D. Dentcheva, A. Ruszczyński. 2009. *Lectures on Stochastic Programming: Modeling and Theory*. SIAM, Philadelphia.
- Smola, A. J., S. V. N. Vishwanathan, T. Hofmann. 2005. Kernel methods for missing variables. *Proc. 10th Internat. Workshop on Artificial Intelligence Statist.*, the Society for Artificial Intelligence and Statistics, Barbados, 325–332.
- Zangwill, W. I. 1969. *Nonlinear Programming: A Unified Approach*. Prentice Hall, Englewood Cliffs, NJ.